Check List the journal of biodiversity data

NOTES ON GEOGRAPHIC DISTRIBUTION

Check List 17 (6): 1659–1669
https://doi.org/10.15560/17.6.1659

PENSOFT

Checklist of free-living microturbellarian flatworms (Turbellaria, Platyhelminthes) in the Suquía River Basin, Córdoba, Argentina

Mariana L. Adami[1,2*], Cristina Damborenea[1,2]

1 División Zoología Invertebrados, Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, La Plata, Argentina • MLA: madami@fcnym.unlp.edu.ar https://orcid.org/0000-0002-6503-9825 • CD: cdambor@fcnym.unlp.edu.ar https://orcid.org/0000-0002-6411-1282

2 CONICET, Argentina

* Corresponding author

Abstract

Free-living microturbellarians are a part of lentic and lotic aquatic assemblages and play an important but underestimated role in aquatic ecosystems. Consequently, they are not included in studies on the dynamics of freshwater aquatic invertebrate communities. We report eight new records of microturbellarians from the Suquía River (Córdoba, Argentina): *Catenula lemnae* Dugès, 1832, *Stenostomum arevaloi* Gieysztor, 1931, *Stenostomum* aff. *amphotum* Marcus, 1945, *Macrostomum* aff. *quiritium* Beklemischev, 1951, *Macrostomum platensis* Adami, Damborenea & Ronderos, 2012, *Microstomum* sp., and *Mesostoma erhenbergii* (Focke, 1836). We discuss these results within the geomorphological history of the river basin.

Keywords

Aquatic invertebrates, microturbellarians, first records, Neotropical region, endorheic system

Academic editor: Fernando Carbayo | Received 13 June 2021 | Accepted 23 November 2021 | Published 10 December 2021

Citation: Adami ML, Damborenea C (2021) Checklist of free-living microturbellarian flatworms (Turbellaria, Platyhelminthes) in the Suquía River Basin, Córdoba, Argentina. Check List 17 (6): 1659–1669. https://doi.org/10.15560/17.6.1659

Introduction

The Suquía River is a closed basin and one of the five main rivers in Córdoba Province, Argentina (Díaz et al. 2018). It flows through an interesting region in which geological and geomorphological changes during the Quaternary had considerable impacts on the current drainage network (Mon and Gutiérrez 2009). The Suquía River belongs to an endorheic system, and evidence suggests that its course was diverted to drain into the Mar Chiquita Lake after geomorphological and tectonic events during the Middle Pleistocene. Prior to these events, these waters flowed northeast to Salado River, a tributary of the Plata Basin (Castellanos 1958; Mon and Gutiérrez 2009).

Córdoba Province is in the Neotropical region, where many studies have reported on the high species diversity of turbellarians in a variety of aquatic environments (e.g., Marcus 1945a, 1945b, 1946, 1948, 1950, 1952; Noreña et al. 2005; Martínez-Aquino et al. 2014; Braccini et al. 2016; Reyes and Brusa 2017).

In Argentina, exhaustive studies have been conducted on the ecological and taxonomic traits of "Turbellaria" (Noreña-Janssen 1995a, 1995b, Brusa et al. 2003, 2008; Noreña et al. 2004; Brusa 2006; Damborenea et al. 2007, 2011; Adami et al. 2012; Lachowich et al. 2016; Adami and Damborenea 2020). Most of these studies were undertaken in the Paraná and Río de la Plata Basin,

©The authors. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

but extensive regions remain unexplored, such as in central Argentina where the turbellarian fauna is unknown.

Freshwater turbellarians can be found in headwater streams associated with large rivers and in wetlands associated with large lakes, and they are part of the bottom, surface, and interstitial benthos (Schwartz and Hebert 1986; Blaustein and Dumont 1990). They can also be found in very small streams and brooks, and on various substrates (e.g., moss, vascular plants, wood snags, detritus, gravel, and sand) in the littoral areas of rivers (Noreña et al. 2005). However, most of ecological surveys of aquatic environments underestimate turbellarians, even though they present high diversity and abundance and play an important role in structuring the ecological community and recycling nutrients to higher trophic levels (Kolasa and Tyler 2010). There are two main reasons for such underestimation. First, there is a lack of trained taxonomists, and second, these organisms must first be identified *in vivo*, so samples must be analyzed shortly after collection.

Considering the above, only three aquatic turbellarians—*Girardia anceps* (Kent, 1930), *Phaenocora unipunctata* (Ørsted, 1843), and *Macrostomum tuba* Graff, 1882—have been reported to date in Córdoba Province (Kawakatsu and Rovasio 1992; Noreña-Janssen 1995b; Adami et al. 2017).

Studies of flora and fauna have been conducted in the Suquía River Basin to generate its biodiversity profile (Hued and Bistoni 2007; Bistoni et al. 2018), but turbellarians remain neglected. Here, we report the free-living microturbellarians recently found in the Suquía River Basin and discuss the outcomes in the light of the evolution and geological history of the current drainage network.

Methods

Data were collected in the Suquía River Basin, Córdoba Province, Argentina (Fig. 1A, B). The basin is formed by the confluence of the San Antonio and Cosquín rivers and receives smaller contributions from Los Chorrillos



Figure 1. Study area. **A.** Red square shows the study area in northern Córdoba province. **B.** Detail of the study area (extracted from Santiago et al. 2018). **C.** Six sampling localities (map data ©2020 Google). La Quebrada Reservoir (yellow circle, site 1; orange circle, site 2; green circle, site 3; red circle, site 4), Colanchanga Stream (light blue circle), Isquitipe (Ceballos) River (dark blue circle).

and Las Mojarras streams. Downstream it receives the waters of numerous streams. The basin covers approximately 7,700 km^2 and has an average annual rainfall of 780 mm (Colladón 2014). The average water temperature ranges from 15.8 to 23.1 °C (Merlo et al. 2017). The Suquía hydrological system consists of three sections. The first section is the upper basin in the highlands of the Sierras Grandes at 1,900 m a.s.l., with headwaters and torrential rivers flowing into the San Roque Reservoir. The second section is the middle basin draining the eastern slopes of the Sierras Chicas, together with the drainage area of Córdoba City; the basin receives the runoff from this city of circa 1.5 million people and other non-point sources of pollution. The third section is the lower basin, downstream from Córdoba City. The river meanders through fields with intensive agricultural activity until it ends in the hypersaline Mar Chiquita Lake at 70 m a.s.l. (Bistoni et al. 2018). Our sampling area is in the middle basin of the river, which includes the provincial nature reserve area La Quebrada Park Natural Water Reserve (Reserva Hídrica Natural Parque La Quebrada in Spanish; Fig. 1C, Table 1).

Biogeographically, the Suquía River endorheic system is included in the Pampean Province (López at al. 2008).

Sampling was performed at six sites, covering a total estimated surface area of 45 km^2 (Table 1). The survey was conducted from 2015 to 2019 during different seasons (Table 2).

A 125-µm-mesh net was used to take three 0.5 L of concentrate samples of filtered water from vegetated areas. The captured specimens were transported to the laboratory and observed alive under a dissecting microscope. Microturbellarians were extracted, examined, and photographed *in vivo* as whole squash mounts under a compound microscope.

Voucher specimens of the examined material were deposited in the Invertebrate Collection of Museo de La Plata, Argentina (MLP), except for *Stenostomum arevaloi*. The few specimens that we collected of this species collapsed during observation and no specimens were vouchered.

Localities referenced herein were taken from the cited literature and the website by Tyler et al. (2006–2021). The maps were prepared using Google Earth (Fig. 1C) and the OMS+relief shading map (Figs. 3, 5, 7, https://www.gpsvisualizer.com/draw/).

Results

Order Catenulida Meixner, 1924
Family Catenulidae Graff, 1905
Genus *Catenula* Dugès, 1832

***Catenula lemnae* Dugès, 1832**
Figures 2A, B, 3A

New records. ARGENTINA – **Córdoba** • Isquitipe River; 31°08′56″S, 064°19′48″W; 1.V. 2019; Mariana Adami leg.; all individuals studied alive; 10 specimens preserved in 100% ethanol, MLP-He 7728.

Identification. Identification based on habitus and morphology, described by Marcus (1945b) and Noreña-Janssen (1995b).

Table 1. Sampling sites, coordinates and brief description about these six sites.

Collecting site	Coordinates	Description
La Quebrada, Site 1	31°09′05″S, 064°21′08″W	Artificial Reservoir (ca. 30 ha)
La Quebrada, Site 2	31°09′02″S, 064°20′43″W	Artificial Reservoir (ca. 30 ha) with dominance of *Azolla* sp. aquatic fern
La Quebrada, Site 3	31°09′10″S, 064°20′31″W	Artificial Reservoir (ca. 30 ha)
La Quebrada, Site 4	31°08′48″S, 064°20′40″W	Artificial Reservoir (ca. 30 ha) with dominance of *Azolla* sp. aquatic fern
Colanchanga Stream	31°08′28″S, 064°21′24″W	Natural stream flowing into the La Quebrada Dike, with dominance of the hydrophyte *Hydrocotyle* sp.
Isquitipe (Ceballos) River	31°08′56″S, 064°19′48″W	Natural stream downstream from La Quebrada Dike near Río Ceballos city, with dominance of the hydrophyte *Hydrocotyle* sp.

Table 2. List of microturbellarian species recorded from the Suquía River. Abundances at different sampling sites and seasons during the survey.

Taxon	Isquitipe (Ceballos) River			La Quebrada Reservoir						Colanchanga
				Site 4	Site 1	Site 2	Site 3	Site 4	Site 3	
	Summer 2015	Winter 2017	Autumn 2019	Winter 2015	Winter 2019	Winter 2019	Winter 2019	Winter 2019	Summer 2019	Winter 2019
CATENULIDA										
Catenula lemnae		11	10	20						
Stenostomum arevaloi					5					
Stenostomum aff. *amphotum*	1		30							
MACROSTOMIDA										
Macrostomum aff. *quiritium*				1					5	2
Macrostomum platensis				1			3			
Microstomum sp.		1	6	1						
RHABDOCOELA										
Dalyellia callvucurai		6								
Mesostoma ehrenbergii					1	50	3	40	1	

Individuals form chains of 2–5 zooids (Fig. 2A). Body length 1.4 mm, and width 100 µm at the level of the statocyst. Statocyst single, caudal to the brain (Fig. 2B). Prostomium length 150 µm, a preoral furrow surrounds the body. Anterior end of prostomium thimble-shaped. Triangular mouth posterior to the ciliated groove (Fig. 2B). Protonephridium duct central on the longitudinal axis and running along the zooids.

Remarks and distribution. *Catenula lemnae* was found in large numbers. *Catenula lemnae* shows a wide distribution range (Luther 1960; Marcus 1945b) and has previously been reported in the Paraná River and Río de la La Plata estuary, Argentina (Noreña-Janssen 1995b; Brusa 2005). Thus, this record represents a western distribution range extension to central Argentina of about 500 km.

Family Stenostomidae Vejdovsky, 1880
Genus *Stenostomum* Schmidt, 1848

***Stenostomum arevaloi* Gieysztor, 1931**
Figures 2C–E, 3B

New records. ARGENTINA – **Córdoba** • La Quebrada Reservoir, site 1; 31°09′05″S, 064°21′08″W; 1.VIII.2019; Mariana Adami leg.; 5 individuals studied alive; no voucher specimens available.

Identification. Identification based on habitus and morphology, as described by Marcus (1945b) and Noreña et al. (2005). Body length 1.1 mm and diameter 230 µm in studied specimens. Chains of two zooids (1–1.5 mm). The main morphological features of this taxon are: three pairs of light-refracting bodies associated with both anterior and posterior brain lobes; two pairs of light-refracting bodies above anterior lobes, and one pair of light-refracting bodies above the posterior lobes of brain, this pair frequently formed by two spheres. Pharyngeal glands opening onto the last two-thirds. Anterior region of the intestine with fine dark granules. Simple tail (one appendage) on the dorsal side. Refractive epidermal inclusions present (Fig. 2C–E).

Remarks and distribution. *Stenostomum arevaloi* has a wide distribution both in the Northern Hemisphere (Gieysztor 1931; Nuttycombe and Waters 1938; Kolasa 1973; Lanfranchi and Papi 1978; Kolasa 1991; Larsson and Willems 2010) and in the Southern Hemisphere, but with only one previous report in the Neotropical region (Marcus 1945b; Young and Kolasa 1974). This species was originally described from brackish water.

***Stenostomum* aff. *amphotum* Marcus, 1945**
Figures 4A, B, 5

New records. ARGENTINA – **Córdoba** • Isquitipe River; 31°08′56″S, 064°19′48″W; 8.II.2015; 1.V.2019; Mariana Adami leg.; 4 specimens preserved in 100% ethanol, MLP-He 7729.

Identification. Identification based on habitus and morphology. The specimens agree with the previous



Figure 2. Specimens *in vivo* after squeeze preparation (**A, C, E**) and freehand drawings (**B, D**). **A, B.** *Catenula lemnae*. **C–E.** *Stenostomum arevaloi*. Abbreviations: cg = ciliated groove; cp = ciliated pits; ei = epidermal inclusion; ig = intestinal granules; m = mouth; ph = pharynx; pr = protonephridium; rb = light-refracting body; st = statocyst; t = tail simple.

descriptions (Marcus 1945a, 1945b; Noreña et al. 2005; Damborenea et al. 2011). Body length 0.8 mm and diameter 200 µm. Dorso-lateral ciliated pits and anterior end slightly tapering, with a small constriction at the level of the oral pore. Two light-refracting bodies, with spherical corpuscles associated with the posterior cerebral lobes. Oral pore oval. Pharynx length 150 µm with two types of pharyngeal glands: (1) antero-ventral rounded glands, and (2) groups of elongated club-shaped glands laterally in the posterior half of the pharynx. Muscular pharyngeal sphincter between pharynx and intestine present (Fig. 4A, B). Protonephridium was not observed.

Remarks and distribution. *Stenostomum* aff. *amphotum* shows higher abundance in autumn than in summer



Figure 3. World distribution reported in the literature (yellow circle) and new record (red square). **A.** *Catenula lemnae*. **B.** *Stenostomum arevaloi*.

(Table 2). *Stenostomum amphotum* has been recorded only in South America: Brazil (Marcus 1945b), Peru (Damborenea et al. 2011), and Buenos Aires Province, Argentina (Noreña et al. 2005). Our record is the first from Córdoba Province and the western-most record of this species in Argentina, about 750 km west from the closest previously known occurrence.

Family Macrostomidae van Beneden, 1870
Genus *Macrostomum* O. Schmidt, 1848

***Macrostomum* aff. *quiritium* Beklemischev, 1951**
Figures 4C–E, 7A

New records. ARGENTINA – **Córdoba** • La Quebrada Reservoir, site 3; 31°09′10″S, 064°20′31″W;



Figure 4. Specimens *in vivo* after squeeze preparation (**A, C, F**), and freehand drawings (**B, D, E, G, H**). **A, B.** *Stenostomum* aff. *amphotum*. **C–E.** *Macrostomum* aff. *quiritium*. **F–H.** *Macrostomum platensis*. Details of the penis stylet are shown in **E**, and **H**. Abbreviations: cp = ciliated pits; e = egg; m = mouth; ov = ovary; pg = pharyngeal glands; ps = pharyngeal sphincter; rb = light-refracting bodies; st = stylet; sv = seminal vesicle; t = testis; vg = vesicula granulorum.

20.XI.2019; Mariana Adami leg.; 1 specimen mounted in polyvinyl-lactophenol; MLP-He 7736 • Colanchanga Stream; 31°08′28″S, 064°21′24″W; 1.VIII.2018; Mariana Adami leg.; 1 specimen mounted in polyvinyl-lactophenol; MLP-He 7731 • La Quebrada Reservoir, site 4; 31°08′48″S, 064°20′40″W; 1.VIII.2015; Mariana Adami leg.; 1 specimen mounted in polyvinyl-lactophenol; MLP-He 7730.

Identification. The morphology and genital system agree with previous observations by Adami et al. (2017) and Reyes et al. (2017). Living mature individuals about 1,600 µm long, with maximum width of 400 µm (Fig. 4C, D). Two black eyes about 13 µm in diameter. Gut with folded edges. Needle-shaped copulatory stylet 95 µm to 110 µm long, smaller than the previously reported for Neotropical region (Adami et al. 2017). Stylet narrowing gradually toward the distal oblique end (Fig. 4D, E). The false seminal vesicle not observed, so the identification provisional, as *M.* aff. *quiritium*.

Remarks and distributions. *Macrostomum* aff. *quiritium* were collected in winter in Colanchanga Stream among hydrophytes (*Hydrocotyle* sp.). *Macrostomum quiritium* have been cited for Russia, Poland, and Switzerland (Schärer et al. 2011) and was recently reported in the Neotropical region (Adami et al. 2017). In contrast to the specimens described here, the Peruvian *M. quiritium* reported by Reyes et al. (2017) are slender.

Macrostomum platensis Adami, Damborenea & Ronderos, 2012

Figures 4F–H, 5

New records. ARGENTINA – **Córdoba** • La Quebrada Reservoir, site 3; 31°09′10″S, 064°20′31″W; 20.XI.2019; Mariana Adami leg.; 1 specimen mounted in polyvinyl-lactophenol and 5 specimens preserved in 100% ethanol for DNA-sequencing; MLP-He 7732.

Identification. Our specimens agree with the previous original descriptions of *M. platensis* (Adami et al. 2012). Body length about 1,100 µm and maximum width 460 µm. Two black eyes in the cerebral region. Rhabdites forming bundles of about 4–10. Stylet funnel-shaped and curved, 50 µm long in the specimens reported here, proximal end broad (10 µm), narrowing towards the distal beveled opening end (5 µm) (Fig. 4H). Seminal vesicle spherical, with a thick muscular wall, and vesicula granulorum also muscular (Fig. 4H). Paired, lateral testes in the central region of the body. Paired, lateral ovaries, posterior to testes (Fig. 4F, G).

Remarks and distribution. *Macrostomum platensis* were collected during winter. The vegetation of the sample site was dominated by *Azolla* sp. This is the second



Figure 5. Localities of new records (red square) and previous records (circles) for the following Neotropical species: *Stenostomum* aff. *amphotum* (yellow), *Macrostomum platensis* (green), *Dalyellia callvucurai* (blue). All three new records are represented by the red square.

record of *M. platensis* in the Neotropical region. This species was originally described from a rainwater pond in La Plata locality (Adami et al. 2012).

Family Microstomidae Luther, 1907
Genus *Microstomum* Schmidt, 1848

***Microstomum* sp.**
Figure 6A, B

New records. ARGENTINA – **Córdoba** • Isquitipe River; 31°08′56″S, 064°19′48″W; 1.V.2019; Mariana Adami leg.; 4 specimens preserved in 100% ethanol and 1 fixed in formaldehyde-phosphate buffered saline, MLP-He 7733.

Identification. Body length about 1750 µm and maximum width 650 µm. *Microstomum* sp. with 1–4 zooids (Fig. 6B). Color of body brownish. Rounded anterior and posterior ends (Fig. 6A). Pigmented eyespots absent. Mouth opening ventrally at the anterior end of the body. Pharynx can expand enormously. The elasticity of this anterior part of body enables ingestion of large prey, after which it attains large body diameter. Figure 6A shows a larva of a dipteran in the intestine (Fig. 6A). Preoral intestine extends anterior to the pharynx. Nematocysts present (Fig. 6B). The studied specimens were immature.

Remarks and distribution. *Microstomum* sp. were collected during autumn and winter (Table 2). The specimens could not be identified to species because not all diagnostic characters mentioned by Atherton and Jondelius (2019) were distinguished. *Micortomum lineare* (Müller 1773) was recorded for El Tigre, Santa Fe, Argentina (Noreña-Janssen 1995b), but these specimens had small adhesive papillae in the posterior end, a feature not mentioned by Atherton and Jondelius (2019). Considering the above, a distribution map is not included here.



Figure 6. Specimens in vivo after squeeze preparation (**A, C, F**), and freehand drawings (**B, D, E, G**). **A, B.** *Microstomum* sp. **C–E.** *Dalyellia callvucurai*. **F, G.** *Mesostoma ehrenbergii*. Details of the penis of *D. callvucurai* shown in **E**. Abbreviations: em = embryos; ey = eyes; g = glands; i; intestine; m = mouth; ne = nematocysts; ph = pharynx; pi = preoral intestine; sp = spots; st = stylet.

Family Dalyelliidae von Graff, 1905
Genus *Dalyellia* Gieysztor, 1938

***Dalyellia callvucurai* Damborenea, Brusa & Noreña, 2007**
Figure 5, 6C–E

New records. ARGENTINA – **Córdoba** • Isquitipe River; 31°08′56″S, 064°19′48″W; 2.VIII.2017. Mariana Adami leg.; 2 specimens mounted in polyvinyl-lactophenol, MLP-He 7734.

Identification. Our specimens agree with previous original descriptions of *D. callvucurai* (Damborenea et al. 2007). Body length 1,200–1,400 µm and width 650 µm There is a pair of eyes at the level of the anterior end of the pharynx (Fig. 6C, D). Pharynx barrel-like, long, approximately ⅓ body length. Length of intestine exceeding half the body, leaving the last portion of the body intestine-free. Stylet 290–340 µm in length, formed by two handles (anterior branches), two posterior branches, and numerous fine spines. Anterior handles 200 µm long broadening at the base and joining through a transverse bar, and posterior branches prolonged in a very broad, scimitar-shaped spine 86 µm long (Fig. 6E). Fine spines on the distal region to the posterior handles. More than 20 short spines on ventral surface and six spines; dorsally, six longer spines (Fig. 6E).

Remarks and distribution. Length of fixed specimens from our study are shorter than the specimens described previously because the latter were measured alive. *Dalyellia callvucurai* was originally found in temporary vegetated environments at the Azul locality, Buenos Aires Province (Damborenea et al. 2007). This is the second record of this species in the Neotropical region and is the western-most record in Argentina, about 880 km west of the nearest previously known occurrence.

Family Typhloplanidae Graff, 1908
Genus *Mesostoma* Ehrenberg, 1837

***Mesostoma ehrenbergii* (Focke, 1836)**
Figures 6F, G, 7B

New records. ARGENTINA – **Córdoba** • La Quebrada



Figure 7. World distribution reported in the literature (yellow circle), and new record (red square). **A.** *Macrostomum quiritium*. **B.** *Mesostoma ehrenbergii*.

Reservoir, Site 4; 31°08′48″S, 064°20′40″W; 1.VIII.2019; Mariana Adami leg.; 35 specimens preserved in 100% ethanol, MLP-He 7735.

Identification. The specimens agree with the previous descriptions and illustrations by Noreña-Janssen (1995b) and Noreña Janssen et al. (2016). Mature individuals 8–11 mm long. Body shape spindle-like with tail and spatula-like anterior end. Body translucent and beige, and dorsal pigmentation arranged in fine spots. Eyes bean-like well separated (Fig. 6F, G). Pharynx rosulatus central (in the second third of the body) ventrally oriented. Some specimens with developed embryos (Fig. 6F).

Remarks and distribution. *Mesostoma ehrenbergii* has a cosmopolitan distribution (Noreña Janssen et al. 2016)

and was previously recorded for Argentina from Santa Fe, Buenos Aires, and Río Negro provinces (Brusa et al. 2020). This species can be found in the phytal areas of pools, ponds, and lakes. In South America, it is present in high densities during autumn and winter, which contrasts with the European population (Noreña-Janssen 1995b).

Discussion

The number of freshwater microturbellarian flatworms species recorded in the Neotropical region is underestimated and poorly known in comparison to other regions in the world. This reflects past scientific activity. The Neotropical freshwater species were first described by Marcus during the 1940s and 1950s in Brazil (Marcus 1946, 1958; Brusa et al. 2020). From 1990 to the present, the diversity and distribution pattern of the South American species are better known as result of the work of Argentine and Brazilian scientists (Braccini et al. 2016; Brusa et al. 2020).

Here, we extend the distribution of eight microturbellarian species by about 600 km west of their previously known distributions, and our records are the first from Córdoba Province for these species.

We did not find three previously species reported for the province. *Macrostomum tuba* Graff, 1882 was recently reported (Adami et al. 2017), but from an artificial pond near the study area. *Phaenocora unipunctata* (Ørsted, 1843) was observed in ponds out of the Suquía river system. During our sampling for this study, we found in the Isquitipe and Colanchanga rivers several specimens of the genus *Girardia*, but they were not identifiable to species and outside of the scope of this research.

All taxa reported here have been previously reported for the Neotropical region. *Catenula lemnae*, *Stenostomum arevaloi*, *Microstomum* sp., *Mesostoma ehrenbergii*, and *Macrostomum quiritium* are widely distributed, present in both the Southern (Neotropical region: Marcus 1943, 1945a, 1945b, 1946; Ponce de León 1984; Braccini et al. 2016; Adami et al. 2017) and Northern hemispheres (Larsson and Willems 2010; Schärer et al. 2011; Noreña et al. 2016; Atherton and Jondelius 2019). *Mesostoma ehrenbergii* has also been recorded in Australia (Noreña et al. 2016). *Macrostomum quiritium* was reported in the Northern Hemisphere were it has been found in artificial ponds with aquatic vegetation from the tropics (Schärer et al. 2011); it only occurs in natural environments in the Neotropical region (Adami et al. 2017; Reyes and Brusa 2017).

In this study, we recorded *Stenostomum arevaloi*, *M. platensis*, and *D. callvucurai* each for only the second time in the Neotropical region (Marcus 1945b; Damborenea et al. 2007; Adami et al. 2012) and extend their distributions. *Macrostomum platensis*, *D. callvucurai*, and *S. amphotum* have been previously found only in the Neotropical region, and they are probably endemic to this region.

It is not surprising that the taxa reported here show a high similarity to the species from the Plata basin (Paraná River and Río de la Plata estuary), except for *Stenostomum arevaloi*, which is recorded for the first time from Argentina.

The Suquía River belongs to an endorheic system. Evidence suggests that its course was diverted to drain into the Mar Chiquita Lake after geomorphological and tectonic events during the Middle Pleistocene (ca. 500,000 years ago). Prior to these events, these waters flowed northeast to the Plata Basin (Castellanos 1958; Mon and Gutiérrez 2009). Freshwater fish and mollusk fauna provide evidence of this ancient connection (López et al. 2008; Núñez et al. 2010; Bistoni et al. 2018). The geological and environmental history of the hydrological system where organisms live are useful in understanding distribution patterns (Formica et al. 2015; Ribeiro et al. 2016).

Microturbellarian species play important roles in freshwater ecosystems and therefore should not be underestimated in ecological studies of the Suquía River. Their taxonomic identification increases the knowledge of biodiversity of the main basin in Córdoba and lays the foundations for research on ecology of microturbelarians and their interactions with other organisms.

Acknowledgements

We thank the ranger of La Quebrada Park Natural Water Reserve, Lucía Gómez for laboratory work, and Marcela Cioccale for providing the literature on the geological history of Córdoba rivers and for her critical reading of the manuscript. Finally, the original manuscript was greatly improved by comments and criticisms from the editor and reviewers. This work was supported by Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata (N886), and PICT2016-3290.

Authors' Contributions

Investigation: MLA. Funding acquisition: CD. Project administration: MLA, CD. Writing – original draft: MLA. Visualization: MLA, CD. Writing – review and editing MLA, CD.

References

Adami ML, Damborenea C (2020) Life history of the freshwater microturbellarian *Macrostomum velastylum* (Macrostomorpha) in the Neotropical Region. Studies on Neotropical Fauna and Environment. https://doi.org/10.1080/01650521.2020.1835294

Adami M, Damborenea C, Ronderos J (2012) A new limnic species of *Macrostomum* (Platyelminthes: Macrostomida) from Argentina and its muscular arrangement labeled with phalloidin. Zoologischer Anzeiger 251: 197–205. https://doi.org/10.1016/j.jcz.2011.08.006

Adami ML, Brusa F, Ronderos JR, Damborenea C (2017) Muscular pattern in three species of *Macrostomum* (Platyhelminthes, Macrostomorpha). Journal of Morphology 278 (2): 264–282. https://doi.org/10.1002/jmor.20633

Atherton S, Jondelius U (2019) A taxonomic review and revisions of Microstomidae (Platyhelminthes: Macrostomorpha). PLoS ONE 14 (4): e0212073. https://doi.org/ 10.1371/journal.pone.0212073

Bistoni MA, Chiarini F, Mangeaud A, Tatián M, Torres R, Visintín A (2018) Biota along the Suquía River. In: Wunderlin DA (Ed.) The Suquía River Basin (Córdoba, Argentina): an integrated study on its hydrology, pollution, effects on native biota and models to evaluate changes in water quality. Springer International Publishing, Cham, Switzerland, 61–92. https://doi.org/10.1007/698_2016_455

Blaustein L, Dumont HJ (1990) Typhloplanid flatworms (*Mesostoma* and related genera): mechanisms of predation and evidence that they structure aquatic invertebrate communities. Hydrobiologia 198: 61–77.

Braccini JAL, Amaral SV, Leal-Zanchet AM (2016) Microturbellarians (Platyhelminthes and Acoelomorpha) in Brazil: invisible organisms? Brazilian Journal of Biology 76 (2): 476–494. https://doi.org/10.1590/1519-6984.21514

Brusa F (2005) Biodiversidad de turbelarios (Platyhelminthes) de la cuenca del Plata. PhD thesis, Universidad Nacional de La Plata. Facultad de Ciencias Naturales y Museo, La Plata, Argentina, 196 pp.

Brusa F (2006) Macrostomida (Platyhelminthes: Rhabditophora) from Argentina, with description of two new species of *Macrostomum* and of stylet ultrastructure. Zoological Sciences 23: 853–862. https://doi.org/10.2108/zsj.23.853

Brusa F, Damborenea MC, Noreña C (2003) A new species of *Gieysztoria* (Platyhelminthes: Rhabdocoela) from Argentina and a kinship analysis of South American species of the genus. Zoologica Scripta 32: 449–457. https://doi.org/10.1046/j.1463-6409.2003.00126.x

Brusa F, Damborenea MC, Noreña C (2008) "Dalyellioida" (Platyhelminthes: Rhabdocoela) from the Río de la Plata estuary in Argentina, with the description of two new species of *Gieysztoria*. Zootaxa 1861: 1–16. https://doi.org/10.11646/zootaxa.1861.1.1

Brusa F, Leal-Zanchet AM, Noreña C, Damborenea C (2020) Phylum Platyhelminthes. In: Damborenea C, Thorp J, Rogers DC (Eds.) Key to Neotropical and Antarctic fauna: Thorp and Covich's freshwater invertebrates, 4th edition. Academic Press, Elsevier, Boston, USA, 101–119. https://doi.org/10.1016/B978-0-12-804225-0.00005-8

Castellanos A (1958) Posibles desplazamientos morfológicos en el pasado de las redes potamográficas en la llanura cordobesa. Separata del Boletín de Estudios Geográficos, Córdoba 19: 29–63.

Colladón L (2014) Síntesis pluviométrica 1992–2012, Cuenca del río San Antonio: sistema del Río Suquía, Provincia de Córdoba. Instituto Nacional del Agua, Ezeiza, 28 pp.

Damborenea C, Brusa F, Noreña C (2007) New Dalyelliidae (Platyhelminthes, Rhabditophora) from Buenos Aires province, Argentina, and their stylet ultrastructure. Zoological Science 24: 803–810. https://doi.org/10.2108/zsj.24.803

Damborenea C, Brusa F, Almagro I, Noreña C (2011) A new species of *Stenostomum* (Platyhelminthes, Catenulida) from the Peruvian Amazon basin and a phylogenetic analysis of Neotropical congeners. Invertebrates Systematics 25: 155–169. https://doi.org/10.1071/IS10026

Díaz E, Corral M, Lábaque M, Vicario L, Pozzi Piacenza C, Moya G, García CM, Tarrab L, Rodríguez A (2018) Hydrology and hydraulics of the Suquía River Basin. In: Wunderlin DA (Ed.) The Suquía River Basin (Córdoba, Argentina): an integrated study on its hydrology, pollution, effects on native biota and models to evaluate changes in water quality. Springer International Publishing, Cham, Switzerland, 1–36. https://doi.org/10.1007/698_2016_466

Dugés A (1832) Description d'un zoophyte, voisin des Bothriocephales (*Catenula Lemnae* Nob). Annales des Sciences Naturelles 26: 198–205.

Focke (1836) *Planaria ehrenbergii*. Annalen des Naturhistorischen Museums in Wien 1 (2): 190–206.

Formica SM, Sacchi GA, Campodonico VA, Pasquini AI, Cioccale MA (2015) Modelado de calidad de agua en ríos de montaña con impacto antrópico. Caso de estudio: Sierra Chica de Córdoba, Argentina. Revista Internacional de Contaminación Ambiental 31 (4): 327–341.

Gieysztor M (1931) Contribution à la connaissance des turbellariés rhabdocèles d'Espagne. Bulletin international de l'Académie Polonaise des Sciences et des Lettres, Classe des sciences mathématiques et naturelles, Série B, Sciences Naturelles 2: 125–153, pls. 13, 14.

Hued AC, Bistoni M (2007) Abundancia y distribución de la fauna íctica en la cuenca del río Suquía (Córdoba, Argentina) Iheringia 97 (3): 286–292. https://doi.org/10.1590/S0073-47212007000300012

Kawakatsu M, Rovasio RA (1992) Redescription of *Dugesia anceps* (Kenk, 1930) (Turbellaria: Tricladida: Paludicola), from the vicinity of Córdoba, Argentina. Proceedings of the Japanese Society of Systematic Zoology 48: 7–23.

Kolasa J (1973) Turbellaria and Nemertini of greenhouses in Poznan. Acta Hydrobiologica 15: 227–245.

Kolasa J (1991) Flatworms: Turbellaria. In: Thorp JM, Covich AP (Eds.) Ecology and classification of North American freshwater invertebrates. Academic Press, New York, USA, 145–171.

Kolasa J, Tyler S (2010) Chapter 6: Flatworms: Turbellarians and Nemertea. In: Thorp, JH, Covich AP (Eds.) Ecology and classification of North American freshwater invertebrates. Academic Press, Elsevier, London, UK, 143–161. https://doi.org/10.1016/B978-0-12-374855-3.00006-6

Lachowicz MC, Adami M, Damborenea C (2016) Variación espacio-temporal de la diversidad de microturbelarios en las provincias de Buenos Aires y Entre Ríos. In: VII Congreso Argentino de Limnología, Tucumán, Argentina. Acta Zoológica Lilloana 60 (Suplemento): 44.

Lanfranchi A, Papi F (1978) Turbellaria (excl. Tricladida). In: Illies J (Ed.) Limnofauna Europaea. 2nd edition. Gustav Fischer Verlag, Stuttgart, Germany, 5–15.

Larsson K, Willems W (2010) Report on freshwater Catenulida (Platyhelminthes) from Sweden with the description of four new species. Zootaxa 2396: 1–18. https://doi.org/10.5281/zenodo.193965

López H, Menni R, Donato M, Miquelarena A (2008) Biogeographical revision of Argentina (Andean and Neotropical regions): an analysis using freshwater fishes. Journal of Biogeography 35: 1564–1579. https://doi.org/10.1111/j.1365-2699.2008.01904.x

Luther A (1960) Die Turbellarien Ostfennoskandiens. I Acoela, Catenulida, Macrostomida, Lecithoepitheliata, Prolecithophora und Proseriata. Fauna Fennica 7: 1–155.

Marcus E (1943) O turbelário *Mesostoma erhenbergii* (Focke 1836) no Brasil. Boletim de Industria Animal (Nova Série) 6: 12–15.

Marcus E (1945a) Sobre Catenulida Brasileiros. Boletim da Faculdade de Filosofia, Ciências e Letras da Universidade de São Paulo, Série Zoologia 10: 3–133.

Marcus E (1945b) Sobre Microturbelarios do Brasil. Comunicaciones Zoológicas del Museo de Historia Natural de Montevideo 1: 1–74.

Marcus E (1946) Sobre Turbellaria brasileiros. Boletim da Faculdade de Filosofia, Ciências e Letras da Universidade de São Paulo, Série Zoologia 11: 1–254.

Marcus E (1948) Turbellaria do Brasil. Boletim da Faculdade de Filosofia, Ciências e Letras da Universidade de São Paulo, Série Zoologia 13: 111–243.

Marcus E (1950) Turbellaria Brasileiros (8). Boletim da Faculdade de Filosofia, Ciências e Letras da Universidade de São Paulo, Série Zoologia 15: 5–192.

Marcus E (1952) Turbellaria Brasileiros (10). Boletim da Faculdade de Filosofia, Ciências e Letras da Universidade de São Paulo, Série Zoologia 17: 5–187.

Marcus E (1958) On South American Turbellaria. Anais dal Academia Brasileira de Ciencias 30: 391–417.

Martínez-Aquino A, Brusa F, Damborenea C (2014) Checklist of freshwater symbiontic temnocephalans (Platyhelminthes: Rhabditophora: Temnocephalida) from the Neotropics. Zoosystematics and Evolution 90: 147–162. https://doi.org/10.3897/zse.90.8688

Merlo C, Abril A, Ame´ MV, Argüello GA, Carreras HA, Chiappero MS et al (2017) Integral assessment of pollution in the Suquía River (Córdoba, Argentina) as a contribution to lotic ecosystem restoration programs. Science of the Total Environment 409 (23): 5034–5045. https://doi.org/10.1016/j.scitotenv.2011.08.037

Mon R, Gutierrez AA (2009) The Mar Chiquita Lake: An indicator of intraplate deformation in the central plain. Geomorphology 111: 111–122. https://doi.org/10.1016/j.geomorph.2009.04.009

Noreña C, Damborenea C, Brusa F (2004) Platyhelminthes de vida libre -Microturbellaria- dulceacuícolas en Argentina. In: F.G. Aceñolaza (Ed.) Temas de la Biodiversidad del Litoral fluvial argentino. INSUGEO, Miscelánea 12: 225–238.

Noreña C, Damborenea C, Brusa F (2005) A taxonomic revision of South American species of the genus *Stenostomum* O. Schmidt (Platyhelminthes: Catenulida) based on morphological characters. Zoological Journal of the Linnean Society 144 (1): 37–58. https://doi.org/10.1111/j.1096-3642.2005.00157.x

Noreña C, Damborenea C, Brusa F (2015) Phylum Platyhelminthes. In: Thorp J, Rogers DC (Eds.) Ecology and general biology, Thorp and Covich's freshwater invertebrates, 4th edition. Academic Press, Elsevier, Boston, USA, 181–203. https://doi.org/10.1016/B978-0-12-385026-3.00010-3

Noreña Janssen C, Damborenea C, Brusa F (2016) Platyhelminthes: Rhabdocoela: Typhloplanidae. In: Dumont HJF (Ed.) Identification guides to the plankton and benthos of inland waters. Margraf Publishers, Weikersheim, Germany, 264 pp.

Noreña-Janssen C (1995a) Studies on the taxonomy and ecology of the Turbellaria (Plathelminthes) in the floodplain of the Paraná River (Argentina). Archiv für Hydrobiologie, Supplement, Algological studies 107: 187–210.

Noreña-Janssen C (1995b) Studies on the taxonomy and ecology of the Turbellaria (Platyhelminthes) in the floodplain of the Paraná River (Argentina). II. Taxonomy and ecology of the Turbellaria. Archiv für Hydrobiologie 107 (2): 211–262.

Núñez V, Gutiérrez-Gregoric DE, Rumi A (2010) Freshwater gastropods provinces from Argentina. Malacologia 53: 47–60. https://doi.org/10.4002/040.053.0103

Nuttycombe J, Waters A (1938) The American species of the genus *Stenostomum*. American Philosophical Society 79: 213–301.

Ponce de León R (1984) Turbellaria del Uruguay II: Sobre *Mesostoma ehrenbergi* (Focke) (Rhabdocoela, Typhloplanidae). Revista de la Facultad de Humanidades y Ciencias. Serie Ciencias Biológicas 1: 381–391.

Reyes J, Brusa F (2017) Species of *Macrostomum* (Macrostomorpha: Macrostomidae) from the coastal region of Lima, Peru, with comments on *M. rostratum* Papi, 1951. Zootaxa 4362: 246–258. https://doi.org/10.11646/zootaxa.4362.2.4

Ribeiro AC, de Lima FCT, Menezes NA (2016) Biogeografia dos peixes de água doce da América do Sul. In: de Carvalho CJB, Almeida EAB (Eds.) Biogeografia da América do Sul. Análise de tempo, espaço e forma. Editora Roca, Río de Janeiro, Brazil, 245–258.

Santiago AN, Bonansea RI, Valdés ME (2018) Organic pollutant in the Suquía River Basin. In: Wunderlin DA (Ed.) The Suquía River Basin (Córdoba, Argentina): an integrated study on its hydrology, pollution, effects on native biota and models to evaluate changes in water quality. Springer International Publishing, Cham, Switzerland, 145–180. https://doi.org/ 10.1007/698_2015_431

Schärer L, Littlewood DTJ, Waeschenbach A, Yoshida W, Vizoso DB (2011) Mating behaviour and the evolution of sperm design. Proceedings of the National Academy of Science of the United States of America 108: 1490–1495. https://doi/10.1073/pnas.1013892108

Schmidt O (1848) Die rhabdocoelen Strudelwürmer (Turbellaria Rhabdocoela) des süssen Wassers. Friedrich Mauke, Jena, Germany, 65 pp., 6 pls.

Schwartz SS, Hebert PDN (1986) Prey preference and utilization by *Mesostoma lingua* (Turbellaria, Rhabdocoela) at a low arctic site. Hydrobiologia 135: 251–253.

Tyler S, Artois T, Schilling S, Hooge M, Bush LF (Eds.) (2006–2021) World list of turbellarian worms: Acoelomorpha, Catenulida, Rhabditophora. http://www.marinespecies.org/turbellarians. Accessed on: 2021-08-05.

Young JO, Kolasa J (1974) Studies on the genus *Stenostomum* O. Schmidt (Turbellaria; Catenulida). Freshwater Biology 4: 167–176.